

# Factors Affecting the Success of Natural Regeneration in Oriental Beech (*Fagus orientalis* Lipsky) Forests in Turkey

Halil Barış ÖZEL<sup>a\*</sup> – Murat ERTEKİN<sup>a</sup> – Mustafa YILMAZ<sup>b</sup> – Erol KIRDAR<sup>a</sup>

<sup>a</sup>Department of Silviculture, University of Bartın, Bartın, Turkey

<sup>b</sup>Department of Silviculture, University of Kahramanmaraş Sütçü İmam, Kahramanmaraş, Turkey

**Abstract** – The success of natural regeneration using the Uniform Shelterwood Method (USM) was determined in an oriental beech (*Fagus orientalis* Lipsky.) stand, in the Bartın-Sökü Forest Range District. Number, height growth and root collar diameter of seedlings were investigated in the 10 experimental plots in the 5.0 ha regeneration area for 7 years (2004–2010). According to the result of factor analyses, it was found that amount of filled seeds, soil cover, growth status of seed trees, stand canopy closure, average crown projection area of seed trees, amount of organic substance in the upper soil layer, soil reaction in the upper soil layer and hillside status were the most effective factors determining the success of natural regeneration.

**Oriental beech (*Fagus orientalis* Lipsky.) / natural regeneration / factor analysis / silviculture**

## 1 INTRODUCTION

Because of variable climate and physiographic conditions, Turkey has natural forest resources with high biological and economic value both with respect to tree species and area. According to 2006 data, forest area of the country is 21.2 million hectares, reaching 27.2% of the total area. 50% of it is degraded and coppice forest. The improvement of this situation and the increase of forestry sector's portion in national income depend largely on successful regeneration practices. Oriental beech (*Fagus orientalis* Lipsky.) has the fourth largest distribution area with 1.7 million ha. Total volume of beech forests is 263.772.103 m<sup>3</sup>, with a total annual increment of 6.130.147 m<sup>3</sup> (OGM 2006).

## 2 MATERIAL AND METHOD

### 2.1 Material

Forests of the Sökü district are situated in the *Northwest Euxinic* forest subzone of the *Euxinic* forest zone (Mayer – Aksoy 1998). According to inventory data for 2000, total forest area in the region is 3386.0 ha, 6.2% (208.5 ha) of it is degraded forest. Sökü region is under the effect of West Black Sea sub-climate (IIc). All seasons are rainy, the month with highest

---

\* Corresponding author: halilbarisozel@yahoo.com; 74100 BARTIN, Turkey

average precipitation is December (162.3mm), the one with the lowest is May (88.0mm). Annual average temperature is 10.6 °C, the coldest month is January (0.0 °C), and the hottest are July (19.8 °C) and August (19.7 °C). Vegetation period in the research area is 6 months (May–October). General soil type is stony, alkaline, sandy clay and sandy clayish mud of mediocre depth (OGM 2001, OGM 2002).

In compartment 27a, a pure beech stand of 5.0 hectares, a regeneration cut using the Uniform Shelterwood Method (USM) has been performed in 2003, in a seed year. During the cut the stand canopy closure has been reduced to 0.6–0.7. Tending of seedlings was continued for 3 years. To follow the increased light demand of seedlings, the canopy closure has been reduced to 0,5 in 2007. A total of 10 experimental plots with the size of 25 x 40 m have been established to follow up the regeneration process.

## 2.2 Method

At first, predominant habitat conditions (climatic, edaphic and physiographic factors such as exposure, slope, altitude and hillside situation) in the experimental plot have been determined. Regarding soil conditions, soil depth, structure type and rooting depth, as well as soil reaction (pH), amount of organic substances, nitrogen, phosphor and potassium have been determined. In addition, in each plot thickness of organic cover (litter and humus layers) and crown cover have been measured. For the seed trees in the 10 experimental plots, age, number, diameter, height, basal area, volume, average annual volume increment, crown width and crown projection area have been determined.

The most important criterion in determining the success of natural regeneration practices is the number of plants per square meter (Saatçioğlu 1979, Atay 1987, Nyland 2002). Not only changes in the number of seedlings but also changes in their height and the root collar diameter per year have been examined. For this purpose, 5 x 5 m sample plots were made up in the 25 x 40 m parcels. In 20 sample plots tally of seedlings, height (cm) and the root collar diameter (mm) measurements were established annually during the seven year research (2004–2010).

## 2.3 Statistical Analysis

SPSS (Statistical Package for Social Science) 9.0 pack program was used for the statistical analysis of data. Kolmogorov-Smirnov test has been applied to determine whether the data displayed normal distribution.

First, correlation analysis has been applied for the purpose of determining the degree of mutual relation between variables and direction. As a result of analysis, dual linear coefficient of correlation between factors have been examined whether they are meaningful or not in the level of significance 0.05 and 0.01. Factor analysis has been applied according to the coefficient of correlation between factors. The Principal Component Analysis has been applied to detect the factors explaining the variance best. “Kaiser” and “Scree Test” criteria are used mostly in determining the number of factors which will represent the relations between variables in the highest degree. In this research, the Kaiser criterion – considering the factors with eigenvalue (joining amount to variance) statistics greater than 1 – was used.


In terms of simplifying the interpretation, unturned factor matrix obtained with Principal Component Analysis should be subjected to rotation. For this purpose, the *Varimax* technique has been applied and a converted factor matrix was obtained by the selection of orthogonal rotation which takes interfactor zero correlation as basis. Interpretation and naming of factors has been made on basis of the variables with high factorial load. In some cases common reasons underlying the variables relating to factors have been taken into consideration. In the research for 7 years, the main aim was to clarify the effects of variables (climatic, edaphic and physiographic and variables relating to seed trees and seedlings) on the number of seedlings.

In this regard, the last three years (2008, 2009 and 2010) data of climate, state of weed and litter cover was used for impact of these variables determined on success of natural regeneration in recent years. For this purpose, multiple regression analysis was carried out in order to determine the effects of all investigated variables on seedling number. The number of beech seedlings per square meter in 2010 has been taken as dependent variable. In multiple regression analysis, the most important factors appeared as independent variables.

### 3 RESULTS AND DISCUSSION

#### 3.1 Number of Seedlings

The success of natural regeneration practices is determined by the regrowth per square meter. Numbers of the last tally carried out in regeneration plots in 2010 is shown in *Figure 1*.




*Figure 1. Mean numbers of oriental beech juvenilities according to experimental areas*

The number of oriental beech seedlings ranged from 2.45 to 12.34 per m<sup>2</sup>. In a study of beech natural regeneration practices in the Belgrade Forest, the average seedling number ranged between 4 and 80 plants/m<sup>2</sup> in the 3rd year and between 4 and 56 plants/m<sup>2</sup> in the 5th year (Saatçioğlu 1970). In Turkish practice, seedling numbers of 2–64 plants/m<sup>2</sup> at the end of 1st year, 1–57 plants/m<sup>2</sup> at the end of 2nd year and 2–71 plants/m<sup>2</sup> at the end of 3rd year are reported (Suner 1978). In the Bartın-Yenihan Forest Range District's 70b compartment, in a study of beech natural regeneration, recruitments between 8 plants/m<sup>2</sup> and 29 plants/m<sup>2</sup> at the end of 1st year, 7 to 28 plants/m<sup>2</sup> at the end of 2nd year and 6 to 26 plants/m<sup>2</sup> at the end of 3rd year were counted (Özel et al. 2009).

When seedling numbers in the present study are compared to other beech natural regeneration practices done with USM, the natural regeneration appears to be unsuccessful. Based on the count in 2010, it looks that 2003 was not a good seed year just a mediocre one in the Bartın-Sökü district. However, one of the reasons why the seedling number is low is that the weed control and soil cultivation in 2003 were not carried out intense enough and with appropriate technique. It was found that the density of weed cover, particularly that of pontic rhododendron (*Rhododendron ponticum* L.) presents an important problem for natural and artificial practices of regeneration in beech. In order to eliminate this problem, clearing of the regeneration area is necessary to remove particularly pontic rhododendron (Saatçioğlu 1970, Kharitonenko 1972, Suner 1978, Eşen 2000).

### 3.2 Height Growth


Average annual height growth of beech seedlings are shown in *Figure 2*. The mean height of 7 years old seedlings, ranged from 14.8 cm to 17.3 cm. In the Belgrade forest, oriental beech seedlings' average height ranged between 13.5 cm and 20.9 cm in the 3rd year, 13.8 cm to 23.0 cm in the 4th and 17.3 cm to 46.6 cm in the 5th (Saatçioğlu 1970). Generally, the height growth of beech seedlings in the Sökü regeneration area is found to be satisfactory.



*Figure 2. Mean heights of oriental beech seedlings in experimental plots*

### 3.3 Root Collar Diameter Development

The root collar diameter growth of seedlings is shown in *Figure 3*. The mean root collar diameter of 7 years old seedlings, ranged from 18.5 mm to 21.3 mm.



*Figure 3. Mean root collar diameter of oriental beech seedlings in experimental plots*

In another research in the Düzce, Cide and Akkuş regions, average root collar diameter after removal cut ranged between 15.8 mm and 23.5 mm (Suner 1978). According to this data, the beech seedlings' root collar diameter growth in the research area is found to be satisfactory too.

### 3.4 Factors Affecting the Regeneration Success

Factor analysis has been applied in order to determine factors that can be effective in the success of regeneration. For this purpose 52 variables were put in the analysis (*Table 1*).

Table 1. Names, dimensions and labels of investigated variables

Order	Variables	Dimension	Label
1	Altitude of Experimental Plot	m	AEA
2	Exposure of Experimental Plot	---	EEA
3	Slope of Experimental Plot	%	SEA
4	Hillside Status of Experimental Plot	---	HSEA
5	Average Diameter of Beech Seed Trees in Experimental Plot	cm	AD
6	Average Height of Beech Seed Trees in Experimental Plot	m	AH
7	Age of Beech Seed Trees in Experimental Plot	Year	ABST
8	Average Crown Form of Beech Seed Trees in Experimental Plot	---	ATF
9	Average Trunk Form of Beech Seed Trees in Experimental Plot	---	ABF
10	Number of of Beech Seed Trees	trees/ha	NBST
11	Average Volume of Beech Seed Trees	m <sup>3</sup> /ha	AV
12	Average Annual Volume Increment of Beech Seed Trees	m <sup>3</sup> /ha/year	AAVI
13	Average Basal Area of Beech Seed Trees	m <sup>2</sup> /ha	ABA
14	Stand Canopy closure	---	SC
15	Mixture Ratio of Beech Seed Trees	%	MR
16	Light Density	%	LI
17	Average Crown Projection Area of Beech Seed Trees in Experimental Plot	m <sup>2</sup>	ATPA
18	Average Crown Width of Beech Seed Trees in Experimental Plot	m	ATW
19	Absolute Soil Depth	cm	ASD
20	Physiological Soil Depth	cm	PSD
21	Rootlet Status	---	RS
22	Structure Type	---	SUT
23	Soil Type	---	ST
24	Soil Reaction in the Upper Soil Layer (pH)	---	SR
25	Amount of Organic Substance in the Upper Soil Layer (Ah)	%	AOS
26	Amount of Nitrogen in the Upper Soil Layer (Ah)	%	AZOTE
27	Amount of Phosphor in the Upper Soil Layer (Ah)	ppm	PHOSPHOR
28	Amount of Potassium in the Upper Soil Layer (Ah)	ppm	POTASSIUM
29	Saltiness of the Upper Soil Layer (Ah)	dS/m	SALTINESS
30	Amount of Filled Seeds	item/ha	AFS
31	Germination Percentage	%	GP
32	Germination Temperature	°C	GT
33	Maturation Time of Seed	day	MTS
34	Damage by Wild Animals	item/m <sup>2</sup>	DWA
35	2008 Amount of Precipitation	mm	AP08
36	2008 Average Temperature	°C	AT08
37	2008 Thickness of Litter Layer	cm	TLL08
38	2008 Thickness of Rotten Layer	cm	TRL08
39	2008 Thickness of Humus Layer	cm	THL08
40	2008 Density of Weed	%	DW08
41	2009 Amount of Precipitation	mm	AP09
42	2009 Average Temperature	°C	AT09
43	2009 Thickness of Litter Layer	cm	TLL09
44	2009 Thickness of Moder (decayed) Layer	cm	TRL09
45	2009 Thickness of Humus Layer	cm	THL09
46	2009 Density of Weed Growth	%	DW09
47	2010 Amount of Precipitation	mm	AP10
48	2010 Average Temperature	°C	AT10
49	2010 Thickness of Litter Layer	cm	TLL10
50	2010 Thickness of Moder (decayed) Layer	cm	TRL10
51	2010 Thickness of Humus Layer	cm	THL10
52	2010 Density of Weed Growth	%	DW10

The number of beech seedlings per square meter in 2010 has been taken as dependent variable and factors affecting this have been examined. In factor analysis, 8 common, basic factors have been identified by using *Principal Component Analysis* and *Kaiser Criterion* (Table 2).

Table 2. Total variance explained for the first 12 components

Components	Initial Eigenvalues			Extraction Sums of Squared Loadings		
	Total	% of Variance	Cumulative Variance (%)	Total	% of Variance	Cumulative Variance (%)
1	28.145	34.312	34.312	15.935	31.853	<b>31.85</b>
2	7.546	16.242	50.554	7.683	20.431	<b>52.28</b>
3	6.845	11.773	62.327	6.815	12.752	65.03
4	3.615	7.665	69.992	5.554	10.026	75.06
5	2.715	5.337	75.329	4.825	8.448	83.51
6	2.413	3.226	78.555	3.673	5.761	89.27
7	2.105	2.475	81.030	2.849	3.275	92.55
8	1.187	1.245	82.275	2.776	1.842	<b>94.39</b>
9	1.170	1.110	83.305			
10	1.163	1.006	84.311			
11	1.152	1.002	85.313			
12	0.996	0.785	86.098			

**Extraction Method:** Principal Component Analysis

As shown in Table 2, first 8 factors with an eigenvalue above 1 are identified. The first factor explains 31.85% of total variance. First and second factors together explain 52.28%. The 8 common, basic factors explain 94.39% of total variance.

In order to facilitate the identification and interpretation the factor, rotated factor matrix has been calculated (Table 3). Factor loads smaller than 0.5 are not included in this chart for the sake of clarity.

Table 3. Rotated component matrix (if more than five components, only the first five shown)

Variables	Factors							
	1	2	3	4	5	6	7	8
AFS	0.999							
GP	0.998							
GT	0.996							
MTS	0.995							
DWA	-0.993							
TLL10		0.994						
TRL10		0.992						
THL10		0.988						
DW10		-0.985						
TLL09		0.982						
AD			0.991					
AH			0.990					
ABST			0.988					
NBST			0.983					
AV			0.980					
SC				0.989				
MR				0.985				
LI				0.980				
ATPA					0.986			
ATW					0.984			
ATF					0.979			
ABF					0.975			
AOS						0.981		
AZOTE						0.980		
PHOSPHOR						0.977		
POTASSIUM						0.973		
ASD						0.970		
SR							0.972	
SALTINESS							-0.967	
HSEA								0.970
AEA								-0.965
EEA								0.953
SEA								-0.932

**Extraction Method:** Principal Component Analysis

**Rotation Method:** Varimax with Kaiser Normalization

As seen in the rotated factor matrix, correlations among 52 variables are represented by 8 common factors. In every factor, the highest correlated variable appears in the first place. Thus, the first variable of the first factor is the amount of filled seeds, the first variable of the second factor is thickness of litter cover in the year 2010; and so on – see *Table 3* and *4*. This way the most influential factors for the success of natural regeneration of oriental beech in the Sökü experiment have been identified. The factors are listed in *Table 4* with their variance contributions.

*Table 4. List of factors most influential in the regeneration success*

The Name of Factor	Variance contribution (%)
1. Amount of Filled Seeds	31.85
2. Soil Cover	20.43
3. Growth Status of Seed Trees	12.75
4. Stand Canopy closure	10.03
5. Average Crown Projection Area of Beech Seed Trees	8.45
6. Amount of Organic Substance in the Upper Soil Layer	5.76
7. Soil Reaction in the Upper Soil Layer	3.28
8. Hillside Status	1.84
Total	94.39

Multivariate regression analysis was performed to examine the effects of the most important nine factors (independent variables) that were identified by the factor analysis on the beech seedlings' number. In the multiple regression analysis, the beech seedlings' number per square meter in the year 2010 (SN.10) was used as dependent variable. Results obtained by Enter Method are given in *Table 5*.

*Table 5. The results of multivariate regression analysis*

Model	Standardized Coefficient	Standard Error	F	R <sup>2</sup>
(Constant)	0.782	0.651		
AFS	0.936**	0.000		
TLL10	0.00453	0.004		
AD	-0.00158	0.003		
SC	0.133**	0.000	7.895***	0.92
ATPA	0.245**	0.000		
AOS	0.237**	0.000		
SR	0.00546	0.007		
HSEA	-0.00047	0.005		

**Dependent variable:** SN.10 (beech seedling number per square meter in the year 2010)

(\*\*): P=0.01 significance level

(\*\*\*): P=0.001 significance level

According to regression analysis result, 92% of the regeneration success in the research area stems from these aforementioned 8 factors. But the most important among them are AFS, SC, ATPA and AOS. These four factors affect seedling number at 99% significance level in the positive direction. According to these results, the multivariate regression model can be written as

$$Y_{(SN10)} = 0.782 + 0.936X_{(AFS)} + 0.00453X_{(TLL10)} - 0.00158X_{(AD)} + 0.133X_{(SC)} + 0.245X_{(ATPA)} + 0.237X_{(AOS)} + 0.00546X_{(SR)} - 0.00047X_{(HSEA)}$$

#### 4 DISCUSSION

Out of the 8 factors that have been obtained as a result of factor analysis, amount of filled seeds takes the first place. There is also an apparent relation between amount of filled seeds and germination percentage of beech seeds. Saatçioğlu (1969), Szwagrzyk et al. (2001) and Çepel (1995) have also found that out of many factors affecting the success of regeneration, the amount of filled seed and germination percentage have evident importances.

The second factor determined by factor analysis is soil cover (*Table 3* and *Table 4*). The soil cover includes weed and litter cover. As the density of weed cover and thickness of litter cover in the regeneration area increases, the number and growth of oriental beech seedlings decreases. It is proven that intense weed cover, particularly pontic rhododendron (*Rhododendron ponticum* L.) cover is an important problem as stated in chapter 3.1 already. Intense weed control and soil cultivation have especially positive effects on seedling development (Saatçioğlu 1970, Suner 1978).

The growth status of seed trees is the third factor having an effect on the success of regeneration (*Table 4*). Trees in the stand should be in the age of maturity in order that regeneration practices can be carried out in a stand (Saatçioğlu 1979, Atay 1987). Trees that haven't reached the full status of maturity yet, cannot yield enough seed, especially as they haven't completed crown growth.

The next factor is the stand canopy closure (*Table 3* and *Table 4*). There is an important relation between stand canopy closure and light density. Light density is effective on germination, site condition and soil cover (weed and litter cover) (Çepel 1995). It is stated by many researchers that oriental beech, growing up slowly in the seedling period, needs protective upper cover until it gains its biological independence (Saatçioğlu 1969, Suner 1978, Atay 1987).

The fifth factor affecting the success of natural regeneration practice in oriental beech is average crown projection area of beech seed trees (*Table 4*). Trees that haven't reached the age of maturity yet, cannot hold enough seed as they haven't completed especially crown development. On the other hand, as age advances, beech spreads its crown and crown decay starts in the upper parts. Because of the decay in the crown of oriental beech trees, capacity of seed holding decreases and this affects the number of seedlings negatively.

The amount of organic substance in the upper soil layer (*Table 3, 4*) is the sixth factor. Seedlings of beech benefit from the water and nutrients in the soil mostly in terms of root growth. In European beech silviculture it is known that the species develops best in soils which are rich in organic substance (Kerr 1995).

The seventh factor is soil reaction in the upper soil layer (*Table 4*). There is a positive relation between amount of soil reaction in the upper soil layer and number of oriental beech seedlings (*Table 3*). In a research carried out by Akgül and Aksoy (1976) in the Research Forest of Bolu-Şerif Yüksel, they confirmed that there is an important relation between root growth, soil reaction and the amount of organic substance.

Hillside status is the eighth factor affecting the success of natural regeneration practice in oriental beech in the research area (*Table 3, 4*). Hillside status is an important physiographic variable influencing local habitat conditions. Depending on the changes in hillside status, microclimatic and edaphic conditions also change dramatically (Çepel 1995). Beech reaches its optimum development at middle and sub hillside (Peters 1992). Outside these conditions the density of beech stand decreases, diameter and height growth decline, crown growth weakens and crown form deteriorates. These changes of the beech trees cause the deterioration of stand structure and fertility.

## 5 CONCLUSIONS

In order to increase the success of beech natural regeneration in the practice the following suggestions are formulated:

- Good seed years should be identified correctly in the stands planned for regeneration.
- The presence of seed trees in sufficient quantity and homogeneity is an important precondition. Regeneration practices should be applied only in stands which are in the age of maturity.
- Before acorn fall in the regeneration area, litter and weed cover, especially pontic rhododendron should be cleared, to provide better conditions for rooting. It should be taken care of mixing litter cover with mineral soil by intensive soil cultivation. After beech seedlings appeared, the necessary tending and weeding should be applied in sufficient frequency and in time.

## REFERENCES

- AGESTAM, E. – EKO, P.M. – NILSSON, U. – WELANDER, N.T. (2003): The effects of shelterwood density and site preparation on natural regeneration of *Fagus sylvatica* L. in southern Sweden, *Forest Ecology and Management* 176: 61–73.
- AKGUL, E. – AKSOY, C. (1976): Bolu-Şerif Yüksel araştırma ormanının toprak karakterleri ve toprak haritaları. [Soil characteristics and soil maps of Bolu Şerif Yüksel Research Forest] OAE, Teknik Bülten No: 95, Ankara, 52 p. (in Turkish)
- ATAY, I. (1971): Tabii Gençleştirmenin Başarılı ve Başarısız Oluşuna Etki Yapan En Önemli Faktörler Üzerine Açıklamalar [The most important factors affecting the success of natural regeneration practices], I.U.Orman Fakültesi Dergisi, B Serisi, Cilt: 2, Sayı: 3: 7–20 İstanbul. (in Turkish)
- ATAY, I. (1987): Doğal Gençleştirme Yöntemleri I-II, [Natural regeneration methods I-II] I.U Fen Bilimleri Enstitüsü, I.U Yayın No: 346, F.B.E Yayın No: 1, İstanbul, 290 p. (in Turkish).
- AVŞAR, M.D. (1999): Kahramanmaraş-Başkonuş Dağı Ormanlarında Başlıca Meşcere Kuruluşları ve Silvikültürel Öneriler. [The main stand structures in the forests of Kahramanmaraş-Başkonuş Mountain and silvicultural suggestions] K.T.Ü. Fen Bilimleri Enstitüsü, Doktora Tezi (yayımlanmamış), Trabzon, 211 p. (in Turkish).
- BOZKUŞ, H.F. (1990): Sedirin (*Cedrus libani* A. Rich.) Toros Göknarı (*Abies cilicica* Carr.) İle Karışık Meşcerelerinde Doğal Gençleştirme Problemleri. [The natural regeneration problems of Lebanon cedar (*Cedrus libani* A. Rich) and Taurus fir (*Abies cilicica* Carr.) mixed stands] OAE Muhtelif Yayınlar No: 59: 435-446, Ankara. (in Turkish)
- ÇEPEL, N. – DUNDAR, M. – GUNEL, A. (1977): Türkiye'nin Önemli Yetiştirme Bölgelerinde Saf Sarıçam Ormanlarının Gelişimi İle Bazı Edafik ve Fizyografik Etkenler Arasındaki İlişkiler. [The relation between pure Scotch pine (*Pinus sylvestris* L.) growth and some edaphic and physiographic factors in important ecological regions in Turkey], Türkiye Bilimsel ve Teknik Araştırma Kurumu (TÜBİTAK), Ankara, 165 p. (in Turkish)
- ÇEPEL, N. (1982): Doğal Gençleştirmenin Ekolojik Koşulları, [Ecological conditions of natural regenerations] I.U Orman Fakültesi Dergisi, B Serisi, Cilt: 32, Sayı: 2: (6–27), İstanbul, (in Turkish).
- ÇEPEL, N. (1995): Orman Ekolojisi. [Forest Ecology], I.U Orman Fakültesi, İstanbul, 536 p. (in Turkish).
- ÇEPEL, N. (2003): Ekolojik Sorunlar ve Çözümleri, [Ecological problems and solutions], Türkiye Bilimsel ve Teknik Araştırma Kurumu (TÜBİTAK), Aydoğdu Matbaası, Ankara, 183 p. (in Turkish)
- ÇOLAK, A.H. (2001): Ormanda Doğa Koruma (Kavramlar, Prensipler, Stratejiler, Önlemler), [Nature conservation in forests (Concepts, Principles, Strategies, Measures)], Orman Bakanlığı, Milli Parklar ve Av-Yaban Hayatı Genel Müdürlüğü, Ankara, 354 p. (in Turkish).

- DAŞDEMİR, İ. (1987): Türkiye'deki Doğu Ladini (*Picea orientalis* L. Carr) Ormanlarında Yetiştirme Ortamı Faktörleri-Verimlilik İlişkisi, [The relation between site factors and productivity in Oriental Spruce (*Picea orientalis* L. Carr) forests in Turkey], İ.Ü. Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi (yayımlanmamış), İstanbul, 122 p. (in Turkish)
- DEMIRCI, A. (1991): Doğu Ladini (*Picea orientalis* (L.) Link.)-Doğu Kayını (*Fagus orientalis* Lipsky.) Karışık Meşcerelerinin Gençleştirilmesi, [Regeneration of oriental spruce (*Picea orientalis* (L.) Link.)-oriental beech (*Fagus orientalis* Lipsky.) mixed stands], K.T.Ü Fen Bilimleri Enstitüsü, Doktora Tezi (yayımlanmamış), Trabzon, 223 p. (in Turkish).
- ELLIOTT, K.J. – KNOEPP, J.D. (2005): The effects of three regeneration methods on plant diversity and soil characteristics in the southern Appalachians, *Forest Ecology and Management* 211: 296-317.
- ERCAN, M. (1997): Bilimsel Araştırmalarda İstatistik, [Statistics in scientific research], Kavak ve Hızlı Gelişen Tür Orman Ağaçları Araştırma Müdürlüğü, İzmit, 225 p. (in Turkish)
- EŞEN, D. (2000): Ecology and Control of Rhododendron (*Rhododendron ponticum* L.) in Turkish Eastern Beech (*Fagus orientalis* Lipsky.) Forests. Faculty of Virginia Polytechnic Institute and State University, Doctora Thesis, Virginia, 111 p.
- GENÇ, M. (2006): Silvikültürel Uygulamalar, [Silvicultural applications] S.D.Ü Orman Fakültesi, Yayın No: 68, Isparta, 357 p. (in Turkish)
- KERR, G. (1995): The silviculture of beech (*Fagus sylvatica* L.) in Europe, genetics and silviculture of beech. In: Proc. 5th Beech Symp. of the IUFRO Project Group P1.10-00, Denmark, 247–256.
- KHARITONENKO, B.Y. (1972): Features of the regeneration of beech in forests of the Black Sea coast of the Caucasus, *Scand. J. For. Res.* 9 (5): 21–23.
- LONG, J. N. – DEAN, T.J. – ROBERTS, S. D. (2004): Linkages between silviculture and ecology: examination of several important conceptual models, *Forest Ecology and Management* 200: 249–261.
- MADSEN, P. – LARSEN, J.B. (1997): Natural regeneration of beech (*Fagus sylvatica* L.) with respect to canopy density, soil moisture, and soil carbon content, *Forest Ecology and Management* 97: 95–105.
- MAYER, H. – AKSOY, H. (1998): Türkiye Ormanları, [Forests in Turkey] Orman Bakanlığı, Batı Karadeniz Ormancılık araştırma Enstitüsü Müdürlüğü, Muhtelif Yayın no:1, Bolu, 291 p. (in Turkish)
- NYLAND, R.D. (2002): *Silviculture (Concepts and applications)*, The McGraw-Hill Company, New York, 682 p.
- OGM (2001): Sökü Orman İşletme Şefliği Model Amenajman Planı, [Model management plan of Sökü Forest Range District] Ankara, 447 p. (in Turkish)
- OGM (2002): Sökü Orman İşletme Şefliği Silvikültür Planı, [Silvicultural plan of Sökü Forest Range District] Bartın, 30 p. (in Turkish)
- OGM (2006): Orman Varlığımız, [State of Turkey's forests] Çevre ve Orman Bakanlığı, Orman Genel Müdürlüğü, OGM Matbaası, Ankara, 152 p. (in Turkish)
- OLIVER, C.D. – LARSON, B.C. (1996): *Forest stand dynamics*, Updated edition, John Wiley & Sons, New York, 520 p.
- ÖZEL, H.B. – ERTEKİN, M. – ÖZDAĞ, S. (2009): Doğu Kayını (*Fagus orientalis* Lipsky.) Ormanlarında Doğal Gençleştirme Başarısını Etkileyen Faktörlerin Belirlenmesi (Bartın-Yenihan Örnek Çalışması), [Determination of factors affecting the success of natural regeneration in oriental beech (*Fagus orientalis* Lipsky.) forests (Bartın-Yenihan case study)], I. Ulusal Batı Karadeniz Ormancılık Kongresi, Cilt: II, Bartın, 471–485. (in Turkish)
- PETERS, R. (1992): *Ecology of beech forests in the Northern Hemisphere*, Wageningen, Netherlands, 125 p.
- PIUSSI, P. – FARRELL, E.P. (2000): Interactions between society and forest ecosystems: challenges for the near future, *Forest Ecology and Management* 132: 21–28.
- SAATÇIOĞLU, F. (1969): *Silvikültürün Biyolojik Esasları ve Prensipleri*, [Biological basics and principles of silviculture], İ.Ü Orman Fakültesi, 1429/138, İstanbul, 323 p. (in Turkish).
- SAATÇIOĞLU, F. (1970): Belgrad Ormanında Kayının (*Fagus orientalis* Lipsky.) Büyük Maktalı Siper Metodu İle Tabii Olarak Gençleştirilmesi Üzerine Yapılan Deney ve Araştırmaların 10 yıllık (1959-1969) Sonuçları, [Ten year results of natural regeneration of oriental beech (*Fagus*

- orientalis* Lipsky.) by using uniform shelterwood methods in Belgrad Forests in Turkey] İ.Ü Orman Fakültesi Dergisi, Cilt: 20, Sayı: 2/A, İstanbul, 1–54. (in Turkish)
- SAATÇIOĞLU, F. (1979): Silvikültür II (Silvikültürün Tekniği), [Silviculture II (Techniques of silviculture)] İ.Ü Orman Fakültesi, 1648/172, İstanbul, 562 p. (in Turkish)
- SCHNEFFER, F. – SCHACHTSCHABEL, P. (2001): Toprak Bilimi (Çevirenler: H. Özbek, Z. Kaya, M. Gök, H. Kaptan), [Soil science] Ç.Ü Ziraat Fakültesi, 73/A-16, Adana, 816 p. (in Turkish)
- SUNER, A. (1978): Düzce, Cide ve Akkuş Mıntıklarında Saf Doğu Kayını Meşcerelerinin Doğal Gençleştirme Sorunları Üzerine Araştırmalar, [Investigation of natural regeneration problems of pure oriental beech stands in Düzce, Cide and Akkuş region in Turkey] Ormancılık Araştırma Enstitüsü, Teknik Bülten Serisi No: 107, Ankara, 60 p. (in Turkish).
- SZWAGRZYK, J. – SZEWCZYK, J. – BODZIARCZYK, J. (2001): Dynamics of seedling bank in beech forest: results of a 10-year study on germination, growth and survival. *Forest Ecology and Management*, 141: 237–250.
- ÜRGENÇ, S. (1998): Ağaçlandırma Tekniği, [Technique of afforestation] İ.Ü Orman Fakültesi, 3994/441, Emek Matbaacılık, İstanbul, 600 p. (in Turkish)

